

Entanglement entropy in aperiodic singlet phases

Róbert Juhász¹ and Zoltán Zimborás²

¹ Research Institute for Solid State Physics and Optics, H-1525 Budapest, P.O.Box 49, Hungary

E-mail: juhasz@szfki.hu

² Research Institute for Particle and Nuclear Physics, H-1525 Budapest, P.O.Box 49, Hungary

E-mail: zimboras@rmki.kfki.hu

Abstract. We study the average entanglement entropy of blocks of contiguous spins in aperiodic XXZ chains which possess an aperiodic singlet phase at least in a certain limit of the coupling ratios. In this phase, where the ground state constructed by a real space renormalization group method, consists (asymptotically) of independent singlet pairs, the average entanglement entropy is found to be a piecewise linear function of the block size. The enveloping curve of this function is growing logarithmically with the block size, with an effective central charge in front of the logarithm which is characteristic for the underlying aperiodic sequence. The aperiodic sequence producing the largest effective central charge is identified, and the latter is found to exceed the central charge of the corresponding homogeneous model. For marginal aperiodic modulations, numerical investigations performed for the XX model show a logarithmic dependence, as well, with an effective central charge varying continuously with the coupling ratio.

PACS numbers: 03.67.Mn, 75.50.Kj

Submitted to: *Journal of Statistical Mechanics: Theory and Experiment*

1. Introduction

The entanglement properties of strongly correlated quantum systems is presently attracting great attention in condensed matter physics [1, 2, 3, 4], quantum information theory [5], and DMRG theory [6]. From the point of view of condensed matter physics, one main aim is to understand the role of quantum correlations in quantum critical phenomena. A widely used estimator of quantum correlations is the *entanglement entropy* of a subsystem s embedded in a larger system, which is defined as

$$S(\hat{\rho}_s) = -\text{Tr } \hat{\rho}_s \ln \hat{\rho}_s , \quad (1)$$

where $\hat{\rho}_s$ denotes the reduced density matrix of the subsystem. Near quantum critical points entanglement entropy was found to display interesting scaling behavior [1, 2, 7, 8]. In one dimensional homogeneous critical spin chains the entanglement entropy belonging to a block of L contiguous spins was shown to grow asymptotically as

$$S_L = \frac{c}{3} \ln L + k , \quad (2)$$

where c is the central charge of the associated conformal field theory, and k is a non-universal constant, while S_L was found to remain bounded in non-critical chains. This type of scaling of the entropy seemed to be related to the conformal invariance of homogeneous critical systems [1, 2]. However, similar logarithmic scaling of the entropy was also found, although, with a different "effective" central charge, in critical inhomogeneous systems such as systems with a single defect [9, 10] or quenched random disorder [3, 11, 12, 13, 14, 15, 16, 17], where the ground state and the low temperature properties are quite different from those of homogeneous models [18]. Some of these recent results have already been used in the experimental identification of certain disordered phases [19].

In the present paper a systematic study of the scaling of the mean entanglement entropy in a wide class of aperiodic models is carried out. The considered models, the self-similarly singlet-renormalizable aperiodic XXZ chains, are characterized by the property that the aperiodic sequence of the couplings remain invariant under a real space RG procedure which produces effective spin singlets. The ground states of these models are aperiodic singlet states, at least in the limit where the RG method applies. Some features of two particular models from this class (the Fibonacci and the tripling chain) have already been investigated in a recent work [20].

We shall derive a general formula for the effective central charge characterizing the asymptotic L -dependence of the average entropy and show that in the XXZ chain this type of inhomogeneity may even enhance entanglement in contrast to single defect and random perturbations, in which cases the effective central charge decreases or remains invariant. Although, it is known that in some special chains of spins with many components also random disorder may increase entanglement [13].

The paper is organized as follows. In Sec.2 some basic features of aperiodic sequences and the renormalization group method for inhomogeneous quantum chains are recapitulated and the class of models to be studied is introduced. In Sec.3 we derive

an analytical expression for the entanglement entropy asymptotics in the considered aperiodic singlet phases, whereas in Sec.4 results of numerical investigations for the XX chain are presented. Sec.5 is devoted to the discussion of the results. Finally, in the Appendix, we prove that among the self-similarly singlet-renormalizable aperiodic XXZ models the one with strong tripling modulation has the largest effective central charge.

2. Aperiodic quantum spin chains

We study the antiferromagnetic spin- $\frac{1}{2}$ XXZ chain defined by the Hamiltonian

$$H = \sum_i J_i (S_i^x S_{i+1}^x + S_i^y S_{i+1}^y + \Delta_0 S_i^z S_{i+1}^z), \quad (3)$$

where the S_i^α 's ($\alpha = x, y, z$) are spin- $\frac{1}{2}$ operators, the site-dependent couplings $J_i > 0$ are taken from a finite set $\{J_a, J_b, \dots\}$ and are modulated according to some aperiodic sequence, whereas Δ_0 ($0 \leq \Delta_0 \leq 1$) is the anisotropy parameter which is initially site-independent, but, as we will see later, it becomes site-dependent in the course of renormalization, as well.

2.1. Aperiodic sequences

The aperiodic sequences considered in this work are composed of letters taken from a finite alphabet $\{a, b, c, \dots\}$ and are defined by an inflation rule which assigns to each letter a word (i.e. a finite sequence of letters) and the (infinitely many) repeated application of which generates the sequence. E.g. the well-known Fibonacci sequence consists of two different letters, a and b , and is defined by the inflation rule

$$\sigma_F : \begin{cases} a \rightarrow w_a = ab \\ b \rightarrow w_b = a. \end{cases} \quad (4)$$

The first few inflation steps starting from letter a results in the following subsequent strings: $a, ab, aba, abaab, abaababa, \dots$. Many properties of aperiodic sequences are encoded in the substitution matrix $M_{\alpha\beta} = n_\alpha(w_\beta)$, where $n_\alpha(w_\beta)$ denotes the number of letters α in the word w_β . E.g. the largest eigenvalue λ_+ of M gives the asymptotic ratio of lengths of the subsequent strings in the inflation procedure.

A finite aperiodic Hamiltonian is constructed by taking a finite section of the infinite aperiodic sequence and assigning a coupling J_α to link $(i, i+1)$ whenever the i th letter of the sequence is α .

2.2. Relevance/irrelevance criteria

According to a heuristic criterion developed originally for random models[21] and later generalized to aperiodic systems[22], the relevance of a weak aperiodic modulation is related to the wandering exponent ω which characterizes the fluctuations of the local order parameter. To be concrete, an aperiodic perturbation is predicted to be relevant if

$$\omega > 1 - \frac{1}{d\nu}, \quad (5)$$

where d is the dimension of the system and ν is the correlation length exponent of the homogeneous system.

For the particular case of aperiodic XX chains an exact relevance/irrelevance criterion is available which is obtained by an exact renormalization group method [23]. Here, the relevant wandering exponent is related to the fluctuations of the non-overlapping pairs of letters, and the aperiodic perturbation is relevant (irrelevant) if $\omega > 0$ ($\omega < 0$), in agreement with the Harris-Luck criterion with $d = 1$ and $\nu = 1$. For $\omega = 0$, which is realized e.g. in the case of Fibonacci modulation, the perturbation is strictly marginal and critical exponents vary continuously with the coupling ratio.

According to a weak perturbation renormalization group study of the general XXZ chain [24] the weak Fibonacci modulation drives the system away from the pure system fixed point for any anisotropy $0 \leq \Delta_0 \leq 1$.

2.3. Strong disorder renormalization group

Recently, many results have been obtained for aperiodic quantum spin chains by means of a real space renormalization group method [25, 26] which was originally developed for random systems [27, 28]. The main point of the method is the successive elimination of high-energy degrees of freedom connected to the largest coupling in the system. For the antiferromagnetic XXZ chain, the spin pair (S_i, S_{i+1}) coupled by the largest bond J_i , which is approximatively in the singlet state $\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ provided that $J_i \gg J_{i-1}, J_{i+1}$, $\Delta_i J_i \gg \Delta_{i-1} J_{i-1}, \Delta_{i+1} J_{i+1}$, is eliminated from the chain, and effective new couplings between spins S_{i-1} and S_{i+2}

$$\tilde{J} = \frac{J_{i-1}J_{i+1}}{(1 + \Delta_i)J_i}, \quad \tilde{\Delta} = \frac{1 + \Delta_i}{2}\Delta_{i-1}\Delta_{i+1} \quad (6)$$

are generated by second order perturbation theory (see Fig. 1). This elimination step




Figure 1. Renormalization scheme in the XXZ chain. The spins connected by the strongest bond J_i form a singlet and a new bond $\tilde{J}$ is generated.

is iterated, leading to the gradual decrease of the number of active spins and that of the energy scale set by the actually largest effective coupling. For the critical random XXZ chain the distribution of couplings is broadening without limits during the procedure, implying that the conditions for the perturbation calculation are better and better fulfilled, and it becomes asymptotically exact in the (infinite randomness) fixed point. The latter is attractive for any amount of disorder, in agreement with the Harris criterion which predicts the random inhomogeneity to be relevant for $0 \leq \Delta_0 \leq 1$. The resulting ground state constructed by the method consists of singlets of spins which are separated by arbitrarily large distances, and the system is in the so called random singlet phase.

The method was applied to the XXZ chain with various types of aperiodic modulations, where identical blocks are simultaneously renormalized in the whole chain. After such a renormalization step the structure of certain aperiodic modulations remain self-similar [25, 26]. E.g. for the Fibonacci XXZ chain which is known to be such a system, the blocks to be renormalized correspond to the words *aba* and *ababa* (see Fig. 2), and the renormalization step is essentially the reversed process of the triple application of σ_F given in eq. (4) [25]. When $\Delta_0 = 1$, the ratio of the couplings




Figure 2. Renormalization of the Fibonacci XXZ chain.

$r_{ab} \equiv J_a/J_b$ and the anisotropy renormalize to $r_{ab}^* = 0$ and $\Delta^* = 1$, respectively. The method becomes asymptotically exact in the fixed point which is attractive for any $r_{ab} < 1$ in agreement with the fact that weak perturbation is relevant. The ground state consists of spin singlets the possible lengths of which form a subset of the Fibonacci numbers and the system is in the so called *aperiodic singlet phase*. When $\Delta_0 = 0$, the ratio r_{ab} remains invariant in the RG procedure, indicating that the Fibonacci modulation is marginal. Here, the decimation steps are exact only in the strong modulation limit $r_{ab} \rightarrow 0$. For the case $0 < \Delta_0 < 1$ the coupling ratio and anisotropy parameter renormalize to some $r_{ab}^* > 0$ and $\Delta^* = 0$, respectively. Thus, the strong perturbation is expected to be marginal whereas the weak perturbation is relevant. Here, the critical exponents are non-universal again, depending both on r_{ab} and Δ_0 [25].

2.4. Singlet producing sequences

In this work, we focus on XXZ chains with such aperiodic sequences (like the Fibonacci chain), where the RG procedure produces exclusively effective singlets and the sequence of couplings remains invariant. Generally, in such systems, the blocks to be renormalized consist of an even number of spins (an odd number of couplings), and the couplings in each block are alternately weak and strong (with strong bonds at even places) in order to avoid the forming of effective spins [26]. Furthermore, not only the RG-invariance of the sequence of couplings (self-similarity) is required, but also the order of strength of the couplings should remain invariant under renormalization, i.e. $J_\alpha > J_\beta$ should imply $\tilde{J}_\alpha > \tilde{J}_\beta$. Each such sequence has a unique *canonical inflation rule* where the word w_α represents that block which just renormalizes to the two-spin block represented by

letter α , that is, the inflation rule just corresponds to the reversed renormalization step. To avoid some unphysical situations we also require two technical conditions: the ergodicity of the inflation rule, i.e. the same infinite sequence should be obtained in the bulk independently of the letter which the iteration is started from, and the RG procedure should be “unambiguous”, i.e. $w_\alpha \neq w_\beta$ if $\alpha \neq \beta$. We call the sequences fulfilling the above conditions *singlet producing self-similar sequences*.

The family of singlet producing self-similar two-letter sequences have the following canonical inflation rule:

$$\sigma_{mn} : \begin{cases} a \rightarrow w_a = aba(ba)^{m-1} \\ b \rightarrow w_b = a(ba)^{n-1}, \end{cases} \quad (7)$$

where the integers m, n satisfy $1 \leq n \leq m$, and letter b now and in the rest of this work represents the strongest bond J_b . The sequence generated by the transformation with $m = n = 1$ is the well-known silver-mean sequence, the choice $m = n = 2$ generates the Fibonacci sequence, whereas the transformation with $m = n = 3$ can be expressed as a composition of simpler inflation transformations as $\sigma_F \circ \sigma_{11} \circ \sigma_F$. An example for a singlet producing sequence composed of three letters is

$$\sigma_{6-3} : \begin{cases} a \rightarrow w_a = abababc \\ b \rightarrow w_b = abc \\ c \rightarrow w_c = ababc, \end{cases} \quad (8)$$

which is the sequence obtained after the first RG step of the 6-3 sequence [26]. Another one is the tripling sequence

$$\sigma_t : \begin{cases} a \rightarrow w_a = aba \\ b \rightarrow w_b = cbc \\ c \rightarrow w_c = abc. \end{cases} \quad (9)$$

which plays a special role among singlet producing sequences as will be shown later. For the latter two sequences the relations $J_b > J_c > J_a$ remain invariant under the renormalization. Generally, the new effective couplings after an RG step are given in terms of the couplings of the previous generation and the canonical substitution matrix as

$$\begin{aligned} \tilde{J}_\alpha &= \frac{\prod_{\gamma \neq b} J_\gamma^{n_\gamma(w_\alpha)}}{J_b^{n_b(w_\alpha)} (1 + \Delta_b)^{n_b(w_\alpha)}}, \\ \tilde{\Delta}_\alpha &= \left(\frac{1 + \Delta_b}{2} \right)^{n_b(w_\alpha)} \prod_{\gamma \neq b} \Delta_\gamma^{n_\gamma(w_\alpha)}, \end{aligned} \quad (10)$$

which are obtained by eliminating the strong bonds J_b one by one in the blocks represented by the words w_α , using eq. (6). The considered sequences can be categorized into three types on the basis of their relevance in the strong perturbation limit.

Those sequences for which the strong perturbation is relevant independently of Δ_0 are of type I. In this case, the coupling ratios renormalize to zero, the procedure is asymptotically exact and the resulting ground state is an aperiodic (independent) singlet state. These aperiodic modulations are exactly known to be relevant in the XX

model for arbitrary coupling ratios, and this is expected to hold also for $0 < \Delta_0 \leq 1$. The two-letter sequences with $m > n$ and e.g. the 6-3 sequence are of type I. Type II sequences, such as the Fibonacci sequence, are relevant for the Heisenberg chain ($\Delta_0 = 1$), but are strictly marginal for the XX chain ($\Delta_0 = 0$). Here, for $\Delta_0 = 0$, the coupling ratios remain invariant under the renormalization, whereas for intermediate initial anisotropies $0 < \Delta_0 < 1$ the coupling ratios tend to a finite limiting value. So, the ground state is an aperiodic singlet state for $\Delta_0 = 1$, but for $\Delta_0 < 1$ this is true only in the limit of vanishing initial coupling ratios. The two-letter sequences with $m = n$ are of this type. Type III modulations are strictly marginal for the XX model, and the coupling ratios remain invariant under renormalization, independently of Δ_0 , thus marginality is indicated (at least in the strong modulation limit). Here, the ground state is an aperiodic singlet state only in the limit of vanishing initial coupling ratios. The tripling sequence and other sequences with words of equal length belong to this class.

3. Entanglement entropy in the aperiodic singlet phase

3.1. Analytical expression for the scaling of entanglement entropy

In this section, we consider an infinite chain and calculate the average entanglement entropy of subsystems consisting of L consecutive spins. In the aperiodic singlet phase this can be done analytically, since, similarly to the random singlet phase, one simply has to count the singlet bonds connecting the subsystem with the rest of the system [3]. The entanglement entropy of one spin in the singlet pair is $\ln 2$, therefore each such bond gives a contribution of $\ln 2$ to the entanglement entropy (Fig. 3). At any stage of




Figure 3. Calculation of the entanglement entropy. In the present case, the entanglement entropy of the block indicated by the arrow is $3 \ln 2$.

the renormalization procedure, the spin pairs, which form singlets, are those connected by the actually strongest (and, at the same time, shortest) bonds, $\tilde{J}_b$. The evolution of bond lengths during the renormalization can be determined by arranging them in a vector w and noticing that the corresponding bond lengths after a renormalization step are obtained by multiplying w by the transpose of the canonical substitution matrix M . The length of the singlets in the n th generation (i.e. those generated in the n th step of the renormalization), which is equal to the length of the bond corresponding to letter b before the n th step, can thus be calculated after diagonalizing M , yielding

$$l_n = u_b \lambda_+^{n-1} + \mathcal{O}(\lambda_-^{n-1}), \quad (11)$$

where λ_+ is the largest eigenvalue of M and u_α denotes the component α of the left eigenvector of M associated with the eigenvalue λ_+ . This eigenvector and the right eigenvector v of M associated with λ_+ are normalized as follows

$$\sum_{\alpha} v_{\alpha} = \sum_{\alpha} u_{\alpha} v_{\alpha} = 1. \quad (12)$$

The terms in the r.h.s. of eq. (11), denoted by $\mathcal{O}(\lambda_+^{n-1})$, involve the subleading eigenvalues of M and are negligible compared to the leading order term for large n . The concentration of singlets of the n th generation, i.e. their average number per unit length, is given by

$$\rho_n = v_b \lambda_+^{-(n-1)}. \quad (13)$$

The fraction of the chain which is covered by singlets of the n th generation is $w_n = l_n \rho_n$, which tends rapidly to $u_b v_b$ for large n . This quantity gives the probability for breaking a singlet bond of the n th generation when the chain is cut at a randomly chosen link. Therefore, each generation for which $l_n < L$ contributes a term of $2w_n \ln 2$ to the average entanglement entropy. The total contribution from singlets with $l_n < L$ is thus for large L

$$S_1(L) \approx 2n(L) u_b v_b \ln 2 + \text{const}, \quad (14)$$

where $n(L)$ is the number of generations with $l_n < L$. From eq. (11) we get for large L

$$n(L) = \left\lfloor \frac{\ln(L/u_b)}{\ln \lambda_+} \right\rfloor_{\text{int}}, \quad (15)$$

where $[x]_{\text{int}}$ stands for the integer part of x . The contribution of singlets with $l_n > L$ is

$$S_2(L) = \sum_{n=n(L)+1}^{\infty} 2\rho_n L \ln 2 = L \lambda_+^{-n(L)-1} \ln 2, \quad (16)$$

where we have used the relation

$$\sum_{n=1}^{\infty} \rho_n = \frac{1}{2}, \quad (17)$$

which expresses the fact that all spins are paired in the aperiodic singlet phase. Finally, we obtain for the asymptotic L -dependence of the entanglement entropy:

$$S(L) = 2n(L) u_b v_b \ln 2 + L \lambda_+^{-n(L)-1} \ln 2 + \text{const}. \quad (18)$$

As opposed to translation invariant systems, $S(L)$ is a piecewise linear function with breaking points at all possible singlet lengths l_n , see Fig. 4. The appearance of the characteristic lengths l_n in the correlation functions is common in the aperiodic singlet phase [26]. Nevertheless, the enveloping curve of $S(L)$, which asymptotically fits to the breaking points, grows logarithmically with L , hence one can write the entropy in the form

$$S(L) = \frac{c_{\text{eff}}}{3} \ln L + P(L), \quad (19)$$

where $P(L)$ is an oscillating log-periodic function. From eqs. (15) and (18) we obtain the general expression for the effective central charge

$$c_{\text{eff}} = 6 \ln 2 \frac{u_b v_b}{\ln \lambda_+}, \quad (20)$$

which differs from the central charge of the homogeneous system $c = 1$ and it is characteristic for the underlying aperiodic sequence. Evaluating eq. (20) for the two-




Figure 4. Average entanglement entropy as a function of the block size L given in eq. (18), evaluated for the Fibonacci sequence (solid curve). The dashed curve is the enveloping function $S_{\text{env}}(L) = \frac{c_{\text{eff}}}{3} \ln L + \text{const.}$

letter sequences σ_{mn} yields

$$c_{\text{eff}} = \frac{6 \ln 2}{\ln \lambda_+} \frac{(\lambda_+ - m - 1)^2}{mn + (\lambda_+ - m - 1)^2}, \quad (21)$$

with the rescaling factor $\lambda_+ = \frac{1}{2}(m + n + \sqrt{(m + n)^2 + 4(m - n + 1)})$. For the 6-3 sequence we get $c_{\text{eff}} = \frac{36 \ln 2}{25 \ln 5} \approx 0.6201$, whereas for the tripling sequence $c_{\text{eff}} = \frac{\ln 4}{\ln 3} \approx 1.2618$.

m=n=1 (silver mean)	0.6910...
m=n=2 (Fibonacci)	0.7962...
m=n=3	0.7819...
m=n=4	0.7519...
m=n=5	0.7228...

Table 1. Effective central charges for aperiodic sequences defined in eq. (7) in the aperiodic singlet phase.




Figure 5. The singlet bond topology of the Fibonacci (upper figure), the silver-mean (middle figure), and the tripling (lower figure) sequences.

3.2. The maximally entangled aperiodic singlet topology

Next, we turn to discuss the possible bounds of the effective central charge. Considering the two-letter sequences, σ_{mn} , for the sake of simplicity with $m = n$, we have found that after the maximum at the Fibonacci sequence ($m = n = 2$) the effective central charge is decreasing with the length of the words (see Table I.) and it finally tends to zero in the limit $n = m \rightarrow \infty$. This is easy to understand, since the words themselves are periodic, and the blocks corresponding to the words have a dimerized (non-critical) ground state. Thus, the effective central charge can be arbitrarily close to zero. One may ask whether sequences (belonging to the considered class) with an arbitrarily large effective central charge can be constructed. The answer is negative. To see this at a heuristic level, note that the expression for the effective central charge in (20) may be separated into two (non-independent) factors, the covered fraction and the rescaling factor. When the singlets of the same generation lie densely, leaving only a small concentration of unpaired spins between them, the covered fraction will be large which is favorable for strong entanglement, however, the rescaling factor will also be large, which has the opposite effect. Conversely, for a large concentration of lonely spins, the rescaling factor is moderate, however, the covered fraction is also small. Thus, one expects that there may be an optimal aperiodic singlet topology which maximizes the entanglement entropy, and indeed this is the case. We can prove that the aperiodic state belonging to the tripling sequence has the highest central charge (see the Appendix). Beside this proof, one can also give a less formal, but more suggestive argument for this. The singlet arrangement produced by the tripling sequence (see Fig. 5) is optimal in the following sense: Regarding those subsequent renormalization steps which generate non-overlapping singlets (e.g. for the silver mean sequence pairs of subsequent steps), as one single renormalization step (in which singlets of different lengths may be produced), the number of unpaired spins between singlets of the same generation is at most one. For the tripling sequence there is an unpaired spin between every neighboring singlet,

which is the optimal situation.

4. Numerical analysis

In order to check the results obtained in the previous section and to get some insight into the behavior of entanglement entropy for marginal perturbations when the ground state is not an independent singlet state and the RG breaks down, we performed numerical calculations for the XX model ($\Delta_0 = 0$). This model can be mapped to a system of free fermions [29] and the reduced density matrix of the subsystem can be written as the exponential of a free-fermion operator [30]. The entanglement entropy can be computed from the eigenvalues λ_i of the correlation matrix of fermion operators $C_{ij} = \langle c_i^\dagger c_j \rangle$ restricted to the subsystem by the formula [1, 30]

$$S(L) = - \sum_i [\lambda_i \ln \lambda_i + (1 - \lambda_i) \ln(1 - \lambda_i)]. \quad (22)$$

The computation of the correlation matrix necessitates the diagonalization of an $N \times N$ matrix, then diagonalizing the restricted correlation matrix of size $L \times L$ one obtains the von Neumann entropy from eq. (22).

We performed numerical calculations for chains with both free and periodic boundary conditions, and calculated the von Neumann entropy of blocks which were located in the former case in the middle of the chain. The size of blocks varied from 2 to $N/2$ and the largest system size we considered was $N = 1024$. The entropies were then averaged over a few thousand independent samples, whereas for periodic chains an averaging over the different starting positions of the block was also carried out.




Figure 6. Numerically calculated average entanglement entropy as a function of the block size L for the relevant two-letter sequence defined in (7) with $m = 2$ and $n = 1$. The calculations were performed with free boundary condition. The size of the system is $N = 1024$ and the data were averaged over 10^4 samples. The solid line is the analytical prediction (18). For this sequence $c_{\text{eff}} \approx 0.4459$.




Figure 7. Numerically calculated average entanglement entropy as a function of the block size L in the Fibonacci XX chain, for different coupling ratios. The calculation was performed with periodic boundary condition. The size of the system is $N = 1024$ and the data were averaged over 20 independent samples and over the different starting positions of the blocks. The solid line is the asymptotics of $S(L)$ for the homogeneous case $r_{ab} = 1$, whereas the dashed curve is the expected asymptotical curve in the limit $r_{ab} \rightarrow 0$, given in eq. (18).

The simplest sequence among the two-letter sequences σ_{mn} which is of type I and is thus relevant even in the XX limit is the sequence with $m = 2$ and $n = 1$. The average entropy as a function of the block size, shown in Fig. 6, tends to the asymptotic curve characteristic for the independent singlet phase after a crossover region even for a finite initial coupling ratio in accordance with the expectations.

For type II sequences, which are marginal in the XX limit and relevant in the Heisenberg limit, one expects that, analogously to the critical exponents, the effective central charge varies with the coupling ratio and the anisotropy parameter for $\Delta_0 < 1$. Numerical results in the XX case for the size dependence of the entanglement entropy for one representant of this class, the Fibonacci sequence, are shown in Fig. 7 for different coupling ratios r_{ab} . As expected, the effective central charge, which is related to the slope of the straight line fitted to the breaking points of the curves in the semi-logarithmic plot, varies continuously with the coupling ratio, see also the numerical results in [20]. For an other type II sequence, the silver-mean sequence, the effective central charge as a function of the coupling ratio is shown in Fig. 8.

Type III sequences, being marginal for any Δ_0 , are also expected to have effective central charges which may depend on the coupling ratios and the anisotropy. Numerical results for the tripling sequence in the XX limit, presented in Fig. 9, are in agreement with these expectations.




Figure 8. Effective central charge for different coupling ratios r_{ab} calculated for the silver-mean XX chain with periodic boundary condition. The arrow indicates the exactly calculated value in the aperiodic singlet phase, taken from Table 1. Inset: The average entropy for different block sizes $L = N/2$ which are chosen to be the first few singlet bond lengths l_n in order to avoid log-periodic oscillations. The averaging was performed on the L different starting positions of the block.




Figure 9. Numerically calculated entanglement entropy as a function of the block size L in the tripling XX chain, for different values of the coupling ratios $J_a/J_c = J_c/J_b \equiv r$. The calculations were performed with free boundary condition. The size of the system is $N = 1024$ and the data were averaged over 10^4 samples. The solid line is the asymptotics of $S(L)$ for the homogeneous case $r = 1$, whereas the dashed curve is the expected asymptotical curve in the limit $r \rightarrow 0$, given in (18).

5. Discussion

We have studied in this work the average entanglement entropy of blocks of contiguous spins in a class of aperiodic quantum spin chains, where the sequence of couplings

remains invariant under the strong disorder RG procedure which produces effective spin singlets. For these systems, the ground state is, in the limit where the RG method applies, an aperiodic (independent) singlet state. Knowing the topology of singlet bonds, the average von Neumann entropy as a function of the block size can be calculated analytically. This quantity, apart from the log-periodic oscillations, is found to grow logarithmically with the size of the block. A general expression for the effective central charge was derived, and the modulation producing the largest central charge was found.

Beside the singlet producing self-similar sequences studied in this work, which can be termed as "pure", also other singlet producing sequences can be constructed by composing canonical inflation transformations of different pure sequences. An example is the transformation $\sigma_{11} \circ \sigma_{33}$ which is, on the other hand, just the square of the transformation $\sigma_{FS} \equiv \sigma_{11} \circ \sigma_F$, therefore both generate the same "golden-silver-mean" sequence. For composite sequences involving m consecutive pure transformations, $M_1, M_2, \dots, M_m$, the calculations outlined in Sec. 3 can be generalized in a straightforward way. In this case the resulting effective central charge is the sum of central charges calculated from the m cyclic permutations of the product $M_1 \circ M_2 \circ \dots \circ M_m$ using eq. (20). For example, for the golden-silver-mean sequence we obtain $c_{\text{eff}} = \frac{6(\sqrt{3}-1)}{\sqrt{3}\ln(7+4\sqrt{3})} \ln 2 \approx 0.6673$.

Finally, we mention that for sequences of type I and II, which are relevant at least for $\Delta_0 = 1$, where the perturbation of arbitrary strength is expected to drive the system to the strongly aperiodic fixed point, we have not found larger values of c_{eff} than that of the homogeneous model $c = 1$. This result is consistent with the "generalized c-theorem" proposed in Ref [3] as a possible extension of the original c-theorem [31] to disordered systems. According to this conjecture, the effective central charge, defined through entanglement scaling, decreases both for homogeneous and disordered systems along RG trajectories induced by relevant perturbations. The sequences for which we have found $c_{\text{eff}} > 1$, the tripling and a sequence with words of length five, both belong to type III. These types of perturbations are marginal at least at the strongly aperiodic fixed point (in the special case $\Delta_0 = 0$ even for perturbations of arbitrary strength), therefore these cases are out of the range of validity of the generalized c-theorem. However, for $\Delta_0 > 0$, the behavior of the effective central charge for finite perturbations of this type is still an open question. It is worth mentioning that our results show that disordered marginal perturbations can increase the effective central charge, which is not expected in case of homogeneous marginal perturbations [32].

Acknowledgments

We thank F. Iglói for useful discussions. This work has been supported by the National Office of Research and Technology under grant No. ASEP1111, by the Hungarian National Research Fund under grant No. T043149 and by the Deutsche Forschungsgemeinschaft under grant No. SA864/2-2.

Appendix A.

Here, we prove that the tripling state has the largest effective central charge among the states generated by singlet producing self-similar sequences.

First, we notice that the relations (13) and (17) imply $v_b = \frac{1}{2}(1 - \lambda_+^{-1})$. Inserting this into the expression of the central charge in eq. (20) we obtain

$$c_{\text{eff}} = 3 \ln 2 \frac{(1 - \lambda_+^{-1})u_b}{\ln \lambda_+}. \quad (\text{A.1})$$

In the considered class of models, as we have mentioned, the strongest effective bond is at the same time the shortest one (or, more precisely, its length is smaller than or equal to that of the other effective bonds) after each renormalization step. Therefore, the length l_{n+1} of singlets produced in the $n + 1$ st renormalization step cannot exceed the average distance ξ_n between active (unpaired) spins after the n th step, i.e. $\xi_n \geq l_{n+1}$. The concentration of active spins decreases exactly by a factor of λ_+ in each step, therefore $\xi_n = \lambda^n$, which, together with eq. (11) leads to $u_b \leq 1$. Putting this in eq. (A.1) yields

$$c_{\text{eff}} \leq 3 \ln 2 \frac{1 - \lambda_+^{-1}}{\ln \lambda_+}. \quad (\text{A.2})$$

Note that we have immediately an upper bound for all the considered sequences, since the r.h.s. of (A.2) is monotonously decreasing with $\lambda_+ (> 1)$, hence $c_{\text{eff}} < 3 \ln 2$. However, this bound can be further sharpened, as we show below.

For $\lambda_+ \geq 3$, the statement we want to prove follows directly, since in this case the maximal value of the upper bound (A.2) is $\frac{\ln 4}{\ln 3}$, which is just the effective central charge of the tripling state.

If $\lambda_+ < 3$, then letter b which represents the strongest bond must be transformed by the canonical inflation rule into a single-letter word. Indeed, $w_b = abc \dots$ would imply $l_{n+1}^b = l_n^a + l_n^b + l_n^c + \dots \geq 3l_n^b$, which is equivalent to $\lambda_+ \geq 3$. Here, $l_n^\alpha = u_\alpha \lambda_+^n + \mathcal{O}(\lambda_+^{-n})$ denotes the length of the bond represented by letter α (or shortly, bond α) after the n th renormalization step. The most general form of canonical inflation rules with $\lambda_+ < 3$, is thus

$$\left\{ \begin{array}{ll} a_1^{(1)} & \rightarrow a_2^{(1)} \\ a_2^{(1)} & \rightarrow a_3^{(1)} \\ & \vdots \\ a_{n_1-1}^{(1)} & \rightarrow a_{n_1}^{(1)} \\ a_{n_1}^{(1)} & \rightarrow w^{(1)} \end{array} \right. \quad \left\{ \begin{array}{ll} a_1^{(2)} & \rightarrow a_2^{(2)} \\ a_2^{(2)} & \rightarrow a_3^{(2)} \\ & \vdots \\ a_{n_2-1}^{(2)} & \rightarrow a_{n_2}^{(2)} \\ a_{n_2}^{(2)} & \rightarrow w^{(2)} \end{array} \right. \quad \vdots$$

$$\left\{ \begin{array}{ll} a_1^{(k)} & \rightarrow a_2^{(k)} \\ a_2^{(k)} & \rightarrow a_3^{(k)} \\ & \vdots \\ a_{n_k-1}^{(k)} & \rightarrow a_{n_k}^{(k)} \\ a_{n_k}^{(k)} & \rightarrow w^{(k)} \end{array} \right.$$

where $a_i^{(j)}$ denotes a single letter, specially $a_1^{(1)} \equiv b$ is the letter representing the strongest bond, whereas $w^{(j)}$ stands for a word comprising at least three letters. The indices n_i are arbitrary positive integers with the restriction $n_1 > 1$. In the following, we use the notation $v_i^{(j)}$ for the density of letter $a_i^{(j)}$ and write $u_i^{(j)} \lambda_+^n$ for the asymptotical length of the corresponding bond after the n th renormalization step. With these notations, eq. (12) reads

$$\sum_{j=1}^k \sum_{i=1}^{n_j} v_i^{(j)} u_i^{(j)} = 1. \quad (\text{A.3})$$

Furthermore, we need the following relations

$$v_{i+1}^{(j)} \geq v_i^{(j)} / \lambda_+, \quad (\text{A.4})$$

$$u_{i+1}^{(j)} = \lambda_+ u_i^{(j)}, \quad (\text{A.5})$$

$$\lambda_+ \geq \sqrt[n]{3}, \quad (\text{A.6})$$

the first two of which follow simply from the form of the inflation rules. Applying eq. (A.5) $n_1 - 1$ times for $j = 1$ yields

$$u_{n_1}^{(1)} = \lambda_+^{n_1-1} u_1^{(1)}. \quad (\text{A.7})$$

Using that $a_{n_1}^{(1)}$ is transformed to the long word $w^{(1)}$ and that $u_1^{(1)} \leq u_i^{(j)}$ for all i, j , we have $\lambda_+ u_{n_1}^{(1)} \geq 3u_1^{(1)}$. This inequality together with eq. (A.7) give immediately relation (A.6).

Now, we consider the group of letters $a_i^{(j)}$ with a fixed index (j) and introduce the quantities $v^{(j)} = \sum_{i=1}^{n_k} v_i^{(j)}$ and $w_i^{(j)} = v_i^{(j)} / v^{(j)}$, such that $\sum_{i=1}^{n_j} w_i^{(j)} = 1$ holds and relations (A.5) imply

$$1 \leq \frac{w_2^{(j)}}{w_1^{(j)}} \lambda_+ \leq \frac{w_3^{(j)}}{w_1^{(j)}} \lambda_+^2 \leq \dots \leq \frac{w_{n_j}^{(j)}}{w_1^{(j)}} \lambda_+^{n_j-1}. \quad (\text{A.8})$$

With these variables we can write

$$\sum_{i=1}^{n_j} v_i^{(j)} u_i^{(j)} = u_1^{(j)} \sum_{i=1}^{n_j} v_i^{(j)} \lambda_+^{i-1} = u_1^{(j)} v^{(j)} \sum_{i=1}^{n_j} w_i^{(j)} \lambda_+^{i-1}, \quad (\text{A.9})$$

where the first equality follows from eq. (A.5). The latter expression can be written also in the form

$$\sum_{i=1}^{n_j} v_i^{(j)} u_i^{(j)} = u_1^{(j)} v^{(j)} \frac{1 + \frac{w_2^{(j)}}{w_1^{(j)}} \lambda_+ + \frac{w_3^{(j)}}{w_1^{(j)}} \lambda_+^2 + \dots + \frac{w_{n_j}^{(j)}}{w_1^{(j)}} \lambda_+^{n_j-1}}{1 + \frac{w_2^{(j)}}{w_1^{(j)}} + \frac{w_3^{(j)}}{w_1^{(j)}} + \dots + \frac{w_{n_j}^{(j)}}{w_1^{(j)}}}. \quad (\text{A.10})$$

One can show that, since $\lambda_+ > 1$, a lower bound of this expression is obtained by substituting the relations (A.8) as equalities into the fraction on the r.h.s. of eq. (A.10), and we get

$$\sum_{i=1}^{n_j} v_i^{(j)} u_i^{(j)} \geq u_1^{(j)} v^{(j)} \frac{n_j \lambda_+^{n_j-1} (\lambda_+ - 1)}{\lambda_+^{n_j} - 1}. \quad (\text{A.11})$$

We know that $u_1^{(j)} \geq u_1^{(1)}$, however, we can sharpen this bound if $n_j < m$, where m is the positive integer for which $\sqrt[m]{3} \leq \lambda_+ < \sqrt[m-1]{3}$. (From (A.6) it follows that $m \geq n_1 > 1$.) In this case, (A.5) yields $u_1^{(j)} \geq 3u_1^{(1)}/\lambda_+^{n_j}$. Using these relations we obtain

$$\sum_{i=1}^{n_j} v_i^{(j)} u_i^{(j)} \geq u_1^{(1)} v^{(j)} P_j(\lambda_+), \quad (\text{A.12})$$

with

$$P_j(\lambda_+) = \begin{cases} \frac{n_j \lambda_+^{n_j-1} (\lambda_+ - 1)}{\lambda_+^{n_j} - 1} & \text{if } n_j \geq m \\ \frac{n_j 3 (\lambda_+ - 1)}{\lambda_+ (\lambda_+^{n_j} - 1)} & \text{if } n_j < m. \end{cases} \quad (\text{A.13})$$

Summing over the index (j) and using the normalization (A.3) we get

$$1 \geq u_1^{(1)} \sum_{j=1}^k v^{(j)} P_j(\lambda_+) \geq u_1^{(1)} \min\{P_j(\lambda_+)\}, \quad (\text{A.14})$$

where the latter inequality follows from $\sum_{j=1}^k v^{(j)} = 1$. It is easy to check that in the interval $\sqrt[m]{3} \leq \lambda_+ < \sqrt[m-1]{3}$

$$\frac{(1 - \lambda_+^{-1})}{P_j(\lambda_+) \ln \lambda_+} \leq \frac{2}{3 \ln 3} \quad (\text{A.15})$$

holds for all $P_j(\lambda_+)$. From this inequality and (A.14) we obtain an upper bound on $u_1^{(1)}$

$$u_b \equiv u_1^{(1)} \leq \frac{2 \ln \lambda_+}{3 \ln 3 (1 - \lambda_+^{-1})}. \quad (\text{A.16})$$

Substituting this into the formula of the effective central charge (A.1) yields

$$c_{\text{eff}} \leq \frac{\ln 4}{\ln 3}. \quad (\text{A.17})$$

The obtained upper bound is again just the value of the effective central charge of the tripling state. Thus, we have completed the proof.

References

- [1] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, Phys. Rev. Lett., **90** 227902 (2003);
J. I. Latorre, E. Rico and G. Vidal, Quant. Inf. and Comp. **4** 048 (2004).
- [2] P. Calabrese and J. Cardy, JSTAT P06002 (2004)
- [3] G. Refael and J.E. Moore, Phys. Rev. Lett., **93** 260602 (2004)
- [4] L. Amico, R. Fazio, A. Osterloh and V. Vedral, preprint, quant-ph/0703044.
- [5] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, U.K., 2000.

- [6] I. Peschel, X. Wang, M. Kaulke, H. Hallberg (Eds.), *Density Matrix Renormalization*, LNP 528, Springer, Berlin, Heidelberg, 1999.
- [7] B.-Q. Jin and V. E. Korepin, J. Stat. Phys. **116**, 79 (2004);
I. Peschel, JSTAT P12005 (2004);
J. P. Keating and F. Mezzadri, Phys. Rev. Lett. **94**, 050501 (2005);
V. Eisler and Z. Zimborás, Phys. Rev. A **71**, 042318 (2005).
Ö. Legeza, J. Sólyom, L. Tincani and R. M. Noack, preprint, cond-mat/0610499.
- [8] M. B. Plenio, J. Eisert, J. Dreissig, M. Cramer Phys. Rev. Lett. **94**, 060503 (2005);
M. Cramer, J. Eisert, M. B. Plenio, and J. Dreissig, Phys. Rev. A **73**, 012309 (2006);
M. M. Wolf, Phys. Rev. Lett. **96**, 010404 (2006);
D. Gioev, I. Klich, Phys. Rev. Lett. **96**, 100503 (2006).
- [9] J. Zhao, I. Peschel, and X. Wang, Phys. Rev. B **73**, 024417 (2006).
- [10] I. Peschel, J. Phys. A **38**, 4327 (2005).
- [11] N. Laflorencie, Phys. Rev. B **72**, 140408(R) (2005).
- [12] G. De Chiara, S. Montangero, P. Calabrese, and R. Fazio, JSTAT P03001 (2006).
- [13] R. Santachiara, JSTAT L06002 (2006).
- [14] J. A. Hoyos and G. Rigolin, Phys. Rev. A **74**, 062324 (2006).
- [15] N. E. Bonesteel and Kun Yang, preprint, cond-mat/0601503.
- [16] A. Saguia, M.S. Sarandy, B. Boechat and M.A. Continentino, preprint, quant-ph/0703028.
- [17] G. Refael and J.E. Moore, preprint, cond-mat/0703038.
- [18] F. Iglói and C. Monthus, Physics Reports **412**, 277 (2005).
- [19] T. G. Rappoport, L. Ghivelder, J. C. Fernandes, R. B. Guimaraes, M. A. Continentino, preprint, cond-mat/0608403.
- [20] F. Iglói, R. Juhász, Z. Zimborás, preprint, cond-mat/0701527.
- [21] A. B. Harris, J. Phys. C **7**, 1671 (1974).
- [22] J. M. Luck, Europhys. Lett. **24**, 359 (1993).
- [23] J. Hermisson, J. Phys. A: Math. Gen. **33**, 57 (2000).
- [24] J. Vidal, D. Mouhanna and T. Giamarchi, Phys. Rev. Lett. **83**, 3908 (1999); Phys. Rev. B **65**, 014201 (2002).
- [25] K. Hida, Phys. Rev. Lett. **93**, 037205 (2004); J. Phys. Soc. Jpn. **73**, 2296 (2004).
- [26] A. Vieira, Phys. Rev. Lett. **94**, 077201 (2005); Phys. Rev. B **71**, 134408 (2005).
- [27] S.-K. Ma, C. Dasgupta, and C.-K. Hu, Phys. Rev. Lett. **43**, 1434 (1979).
- [28] D.S. Fisher, Phys. Rev. B **50**, 3799 (1994).
- [29] E. Lieb, T. Schultz and D. Mattis, Annals of Phys. **16**, 407 (1961).
- [30] I. Peschel, J. Phys. A: Math. Gen. **36**, L205 (2003).
- [31] A. B. Zamolodchikov, JETP Lett. **43**, 730 (1986).
- [32] J. Cardy, J.Phys. A: Math. Gen **20** L891-L896 (1987).